

# Topological equivalences of differential graded algebras

(Joint work with D. Dugger)

---

**“Abelian groups up to homotopy”**

spectra  $\iff$  generalized cohomology theories

**Examples:**

1. **Ordinary cohomology:**

For  $A$  any abelian group,  $H^*(X; A) = [X_+, K(A, n)]$ .

Eilenberg-Mac Lane spectrum, denoted  $HA$ .

$HA_n = K(A, n)$  for  $n \geq 0$ .

The coefficients of the theory are given by

$$HA^*(\text{pt}) = \begin{cases} A & * = 0 \\ 0 & * \neq 0 \end{cases}$$

## 2. Hypercohomology:

For  $C$ . any chain complex of abelian groups,

$$\mathbb{H}^s(X; C.) \cong \bigoplus_{q-p=s} H^p(X; H_q(C.)).$$

Just a direct sum of shifted ordinary cohomologies.

$$HC.*(\text{pt}) = H_*(C.).$$

## 3. Complex K-theory:

$K^*(X)$ ; associated spectrum denoted  $K$ .

$$K_n = \begin{cases} U & n = \text{odd} \\ BU \times \mathbb{Z} & n = \text{even} \end{cases}$$

$$K^*(\text{pt}) = \begin{cases} 0 & * = \text{odd} \\ \mathbb{Z} & * = \text{even} \end{cases}$$

## 4. Stable cohomotopy:

$\pi_S^*(X)$ ; associated spectrum denoted  $\mathbb{S}$ .

$\mathbb{S}_n = S^n$ ,  $\mathbb{S}$  is the *sphere spectrum*.

$\pi_S^*(\text{pt}) = \pi_{-*}^S(\text{pt}) =$  stable homotopy groups of spheres. These are only known in a range.

## “Rings up to homotopy”

ring spectra  $\iff$  gen. coh. theories with a product

1. For  $R$  a ring,  $HR$  is a ring spectrum.

The cup product gives a graded product:

$$HR^p(X) \otimes HR^q(X) \rightarrow HR^{p+q}(X)$$

Induced by  $K(R, p) \wedge K(R, q) \rightarrow K(R, p + q)$ .

**Definition.**  $X \wedge Y = X \times Y / (X \times pt) \cup (pt \times Y)$ .

2. For  $A$ . a differential graded algebra (DGA),

$HA$ . is a ring spectrum. Product induced by

$$\mu : A. \otimes A. \rightarrow A., \text{ or } A_p \otimes A_q \rightarrow A_{p+q}.$$

The groups  $\mathbb{H}(X; A.)$  are still determined by  $H_*(A)$ ,  
but the product structure is *not* determined  $H_*(A)$ .

3.  $K$  is a ring spectrum;


Product induced by tensor product of vector bundles.

4.  $\mathbb{S}$  is a commutative ring spectrum.

## Definitions

**“Definition.”** A *ring spectrum* is a sequence of pointed spaces  $R = (R_0, R_1, \dots, R_n, \dots)$  with compatibly associative and unital products  $R_p \wedge R_q \rightarrow R_{p+q}$ .

**Definition.** The *suspension* of a based space  $X$  is  $\Sigma X = S^1 \wedge X \cong (CX \cup_X CX) / \sim_{\text{pt}}$ .



**Definition.** A *spectrum*  $F$  is a sequence of pointed spaces  $(F_0, F_1, \dots, F_n, \dots)$  with structure maps  $\Sigma F_n \rightarrow F_{n+1}$ .

**Example:**  $\mathbb{S}$  a commutative ring spectrum

Structure maps:  $\Sigma S^n = S^1 \wedge S^n \xrightarrow{\cong} S^{n+1}$ .

Product maps:  $S^p \wedge S^q \xrightarrow{\cong} S^{p+q}$ .

Actually, must be more careful here. For example:  $S^1 \wedge S^1 \xrightarrow{\text{twist}} S^1 \wedge S^1$  is a degree  $-1$  map.

## History of spectra and $\wedge$

*Boardman in 1965 defined spectra and  $\wedge$ .  $\wedge$  is only commutative and associative up to homotopy.*

*$A_\infty$  ring spectrum = best approximation to associative ring spectrum.*

*$E_\infty$  ring spectrum = best approximation to commutative ring spectrum.*

*Lewis in 1991: No good  $\wedge$  exists.*

Five reasonable axioms  $\implies$  no such  $\wedge$ .

*Since 1997, lots of good categories of spectra exist! (with  $\wedge$  that is commutative and associative.)*

1. 1997: Elmendorf, Kriz, Mandell, May

2. 2000: Hovey, S., Smith

3, 4 and 5 ... Lydakis, Schwede, ...

### **Theorem.**

All above models define the same homotopy theory.

## Spectral Algebra

Given the good categories of spectra with  $\wedge$ , one can easily do algebra with spectra.

### Definitions:

A *ring spectrum* is a spectrum  $R$  with an associative and unital multiplication  $\mu : R \wedge R \rightarrow R$  (with unit  $\mathbb{S} \rightarrow R$ ).

An  *$R$ -module spectrum* is a spectrum  $M$  with an associative and unital action  $\alpha : R \wedge M \rightarrow M$ .

$\mathbb{S}$ -*modules* are spectra.

$S^1 \wedge F_n \rightarrow F_{n+1}$  iterated gives  $S^p \wedge F_q \rightarrow F_{p+q}$ .

Fits together to give  $\mathbb{S} \wedge F \rightarrow F$ .

$\mathbb{S}$ -*algebras* are ring spectra.

## Homological Algebra vs. Spectral Algebra

$\mathbb{Z}$	$\mathbb{Z}$ (d.g.)	$\mathbb{S}$
$\mathbb{Z}$ -Mod $= \mathcal{A}b$	d.g.-Mod $= \mathcal{C}h$	$\mathbb{S}$ -Mod $= \mathcal{S}pectra$
$\mathbb{Z}$ -Alg = $\mathcal{R}ings$	d.g.-Alg = $\mathcal{D}GAs$	$\mathbb{S}$ -Alg = $\mathcal{R}ing\ spectra$

$\mathbb{Z}$	$\mathbb{Z}$ (d.g.)	$H\mathbb{Z}$	$\mathbb{S}$
$\mathbb{Z}$ -Mod	d.g.-Mod	$H\mathbb{Z}$ -Mod	$\mathbb{S}$ -Mod
$\mathbb{Z}$ -Alg	d.g.-Alg	$H\mathbb{Z}$ -Alg	$\mathbb{S}$ -Alg

$\mathbb{Z}$	$\mathbb{Z}$ (d.g.)	$H\mathbb{Z}$	$\mathbb{S}$
$\mathbb{Z}$ -Mod	d.g.-Mod	$H\mathbb{Z}$ -Mod	$\mathbb{S}$ -Mod
$\mathbb{Z}$ -Alg	d.g.-Alg	$H\mathbb{Z}$ -Alg	$\mathbb{S}$ -Alg
$\cong$	quasi-iso	weak equiv.	weak equiv.

*Quasi-isomorphisms* are maps which induce isomorphisms in homology.

*Weak equivalences* are maps which induce isomorphisms on the coefficients.

$\mathbb{Z}$	$\mathbb{Z}$ (d.g.)	$H\mathbb{Z}$	$\mathbb{S}$
$\mathbb{Z}$ -Mod	d.g.-Mod	$H\mathbb{Z}$ -Mod	$\mathbb{S}$ -Mod
$\mathbb{Z}$ -Alg	d.g.-Alg	$H\mathbb{Z}$ -Alg	$\mathbb{S}$ -Alg
$\cong$	quasi-iso	weak equiv.	weak equiv.
	$\mathcal{D}(\mathbb{Z}) =$ $Ch[q\text{-iso}]^{-1}$	$\mathcal{H}o(H\mathbb{Z}\text{-Mod})$	$\mathcal{H}o(\mathbb{S}) =$ $Spectra[wk.eq.]^{-1}$

**Theorem.** Columns two and three are equivalent up to homotopy.


- (1) (Robinson '87)  $\mathcal{D}(\mathbb{Z}) \simeq_{\Delta} \mathcal{H}o(H\mathbb{Z}\text{-Mod})$ .
- (2) (Schwede-S.)  $Ch \simeq_{\text{Quillen}} H\mathbb{Z}\text{-Mod}$ .
- (3) (S.)  $DGA \simeq_{\text{Quillen}} H\mathbb{Z}\text{-Alg}$ .
- (4) (S.) For  $A$ . a DGA,  
d.g.  $A$ .-Mod  $\simeq_{\text{Quillen}} HA$ .-Mod  
and  $\mathcal{D}(A) \simeq_{\Delta} \mathcal{H}o(HA$ .-Mod).

rings  $\subseteq$  DGAs  $\subseteq$  Ring Spectra



$$k \rightarrow k[x]$$

Consider DGAs as ring spectra



**Definition.** Two DGAs  $A.$  and  $B.$  are *topologically equivalent* if their associated  $H\mathbb{Z}$ -algebras  $HA.$  and  $HB.$  are equivalent as ring spectra ( $\mathbb{S}$ -algebras).

**Theorem.** If  $A.$  and  $B.$  are topologically equivalent DGAs, then  $\mathcal{D}(A.) \simeq_{\Delta} \mathcal{D}(B.).$

**Proof.** This follows since

$$\begin{aligned} \text{d.g. } A. \text{-Mod} &\simeq_Q HA. \text{-Mod} \simeq_Q HB. \text{-Mod} \\ &\simeq_Q \text{d.g. } B. \text{-Mod} \end{aligned}$$

**Example:**

$$A = \mathbb{Z}[e_1]/(e^4) \text{ with } de = 2 \text{ and } A' = H_*A \\ = \Lambda_{\mathbb{Z}/2}(\alpha_2)$$

degree	A	A' = H_*A
3	$\mathbb{Z} = \langle e^3 \rangle$	
	$\downarrow \cdot 2$	
2	$\mathbb{Z} = \langle e^2 \rangle$	$\mathbb{Z}/2 = \langle \alpha \rangle$
1	$\mathbb{Z} = \langle e \rangle$	
	$\downarrow \cdot 2$	
0	$\mathbb{Z} = \langle 1 \rangle$	$\mathbb{Z}/2 = \langle 1 \rangle$

$A$  and  $A'$  are *not* quasi-isomorphic,  
(although  $H_*A \cong H_*A'$ .)

Claim:  $A$  and  $A'$  are topologically equivalent.  
Or,  $HA \simeq HA'$  as ring spectra.

## Equivalences of module categories

(Morita 1958) Any equivalence of categories  $R\text{-Mod} \cong R'\text{-Mod}$  is given by tensoring with a bimodule.

(Rickard 1989, 1991) Any derived equivalence of rings  $\mathcal{D}(R) \cong_{\Delta} \mathcal{D}(R')$  is given by tensoring with a complex of bimodules (a *tilting complex*).

*also modules  
are Quillen equiv.*

(Schwede-S. 2003) Any Quillen equivalence of module spectra  $R\text{-Mod} \simeq_Q R'\text{-Mod}$  is given by smashing with a bimodule spectrum (a *tilting spectrum*).

(Dugger-S.) Example above shows that for derived equivalences of DGAs one must consider tilting spectra, not just tilting complexes.

In fact, there is also an example of a derived equivalence of DGAs which doesn't come from a tilting spectrum (because it doesn't come from an underlying Quillen equivalence.) (This example is based on work by (Schlichting 2002).)

**Use  $HH^*$  and  $THH^*$ :**

For a ring  $R$  and an  $R$ -bimodule  $M$ , DGAs with non-zero homology  $H_0 = R$  and  $H_n = M$  are classified by  $HH_{\mathbb{Z}}^{n+2}(R; M)$ .

### **Topological Hochschild homology**

Using  $\wedge$  in place of  $\otimes$  one can mimic the definition of  $HH$  for spectra to define  $THH$ .

In particular,  $HH_{\mathbb{Z}}^*(R; M) = THH_{H\mathbb{Z}}^*(HR; HM)$ .

Just as above, ring spectra are classified by  $THH_{\mathbb{S}}^{n+2}(HR; HM)$ .

$A$  and  $A'$  are thus classified in these two settings by letting  $R = \mathbb{Z}/2$ ,  $M = \mathbb{Z}/2$  and  $n = 2$ .

$\mathbb{S} \rightarrow H\mathbb{Z}$  induces

$$\Phi : HH_{\mathbb{Z}}^*(\mathbb{Z}/2; \mathbb{Z}/2) \rightarrow THH_{\mathbb{S}}^*(\mathbb{Z}/2; \mathbb{Z}/2).$$

One can calculate that  $A$  and  $A'$  correspond to different elements in  $HH^4$  which get mapped to the same element in  $THH^4$ .

Compute:

$$HH_{\mathbb{Z}}^*(\mathbb{Z}/2; \mathbb{Z}/2) = \mathbb{Z}/2[\sigma_2]$$

(Franjou, Lannes, and Schwartz 1994)

$$\begin{aligned} THH_{\mathbb{S}}^*(\mathbb{Z}/2; \mathbb{Z}/2) &= \Gamma_{\mathbb{Z}/2}[\tau_2] \\ &\cong \Lambda_{\mathbb{Z}/2}(e_1, e_2, \dots), \quad \deg(e_i) = 2^i. \end{aligned}$$

To compute  $\Phi : HH_{\mathbb{Z}}^*(\mathbb{Z}/2) \rightarrow THH_{\mathbb{S}}^*(\mathbb{Z}/2)$ :

In  $HH^2$ :  $\sigma \leftrightarrow \mathbb{Z}/4$  and  $0 \leftrightarrow \mathbb{Z}/2 \oplus \mathbb{Z}/2$

In  $THH^2$ :  $\tau \leftrightarrow H\mathbb{Z}/4$  and  $0 \leftrightarrow H(\mathbb{Z}/2 \oplus \mathbb{Z}/2)$

So  $\Phi(\sigma) = \tau$ .

In  $HH^4$ :  $\sigma^2 \leftrightarrow A$  and  $0 \leftrightarrow A'$ .

$\Phi(\sigma^2) = \Phi(0) = 0$  since  $\tau^2 = 0$  and  $\Phi$  is a ring homomorphism.

So  $HA \simeq HA'$  as ring spectra,  
although  $A \not\cong A'$  as DGAs.

It follows that  $\mathcal{D}(A) \simeq_{\Delta} \mathcal{D}(A')$ .

**Example:** There exist two DGAs  $A$  and  $B$  such that

$$\mathcal{D}_A \cong_{\Delta} \mathcal{D}_B, \text{ but} \\ \text{d.g.}A\text{-mod} \not\cong_Q \text{d.g.}B\text{-mod}$$

*Based on Marco Schlichting's example ( $p > 3$ ):*

$$\mathcal{H}o(\text{Stmod}(\mathbb{Z}/p[\epsilon]/\epsilon^2)) \cong_{\Delta} \mathcal{H}o(\text{Stmod}(\mathbb{Z}/p^2)), \\ \text{but} \\ \text{Stmod}(\mathbb{Z}/p[\epsilon]/\epsilon^2) \not\cong_Q \text{Stmod}(\mathbb{Z}/p^2)$$

One can find DGAs  $A$  and  $B$  such that:

$$\text{Stmod}(\mathbb{Z}/p[\epsilon]/\epsilon^2) \simeq_Q \text{d.g.}A\text{-mod} \\ \text{Stmod}(\mathbb{Z}/p^2) \simeq_Q \text{d.g.}B\text{-mod}$$

Here  $A$  and  $B$  are the endomorphism DGAs of the Tate resolution of a generator ( $\mathbb{Z}/p$  in both cases):

$$A = \mathbb{Z}/p[x_1, x_1^{-1}] \text{ with } d = 0.$$

$$B = \mathbb{Z}[x_1, x_1^{-1}]\langle e_1 \rangle / e^2 = 0, ex + xe = x^2 \text{ with } de = p \text{ and } dx = 0.$$